

## ERRATUM / ERRATUM

**Erratum: Estimating westslope cutthroat trout (*Oncorhynchus clarkii lewisi*) movements in a river network using strontium isoscapes****Clint C. Muhlfeld, Simon R. Thorrold, Thomas E. McMahon, and Brian Marotz**

**Ref.:** Can. J. Fish. Aquat. Sci. **69**(5): 906–915 (2012).

Figure 4 on page 911 did not show all data. The correct figure is shown on the next page. The publisher apologizes for any inconvenience this might have caused.

Received 1 May 2012. Accepted 1 May 2012. Published at [www.nrcresearchpress.com/cjfas](http://www.nrcresearchpress.com/cjfas) on 7 May 2012.  
J2011-0436E

**C.C. Muhlfeld.** US Geological Survey, Northern Rocky Mountain Science Center, Glacier National Park, West Glacier, MT 59936, USA.

**S.R. Thorrold.** Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA.

**T.E. McMahon.** Department of Ecology, Fish and Wildlife Program, Montana State University, Bozeman, MT 59717, USA.

**B. Marotz.** Montana Fish, Wildlife & Parks, Kalispell, MT 59901, USA.

**Corresponding author:** Clint C. Muhlfeld (e-mail: [cmuhlfeld@usgs.gov](mailto:cmuhlfeld@usgs.gov)).

**Fig. 4.** Mean ( $n = 2$ ) Sr:Ca values (a) and  $^{87}\text{Sr}:^{86}\text{Sr}$  ratios (b) in paired otolith edge and water samples from juvenile cutthroat trout collected in five streams in the North Fork and four streams in the South Fork Flathead River, Montana. Fish from one site (open square) had apparently recently moved to the stream from another location and therefore were omitted from the regression analysis. Solid line represents least squares regression line, shaded area represents 95% confidence intervals for the regression line, and the gray dashed line shows 1:1 relationship between water and otolith values.

